

THE ASYMPTOTIC LEADING TERM FOR MAXIMUM RANK OF TERNARY FORMS OF A GIVEN DEGREE

ALESSANDRO DE PARIS

ABSTRACT. Let $r_{\max}(n, d)$ be the maximum Waring rank for the set of *all* homogeneous polynomials of degree $d > 0$ in n indeterminates with coefficients in an algebraically closed field of characteristic zero. To our knowledge, when $n, d \geq 3$, the value of $r_{\max}(n, d)$ is known only for $(n, d) = (3, 3), (3, 4), (3, 5), (4, 3)$. We prove that $r_{\max}(3, d) = d^2/4 + O(d)$ as a consequence of the upper bound $r_{\max}(3, d) \leq \lfloor (d^2 + 6d + 1)/4 \rfloor$.

Keywords: Waring problem, rank, symmetric tensor.

MSC2010: 15A21, 15A69, 15A72, 14A25, 14N05, 14N15.

1. INTRODUCTION

A natural kind of Waring problem asks for the least of the numbers r such that *every* homogeneous polynomial of degree $d > 0$ in n indeterminates can be written as a sum of r d th powers of linear forms. For instance, when $(n, d) = (3, 4)$ (and the coefficients are taken in an algebraically closed field of characteristic zero), the answer is 7. This was found for the first time in [11]. In view of the interplay with the rank of tensors, relevant applicative interests of questions like this have recently been recognized (see [10]). For further information we refer the reader to [12, Introduction].

Every power sum decomposition gives rise to a set of points in the projectivized space of linear forms, and in [7] it is proved that for ternary quartics one can always obtain a power sum decomposition by considering seven points arranged along three lines. In [8], considering sets of points arranged along four lines, one finds that every ternary quintic is a sum of 10 fifth powers of linear forms. Ternary quintics without power sum decompositions with less than 10 summands were exhibited soon after in [4]. Hence, the answer in the case $(n, d) = (3, 5)$ is 10.

In the present paper we test “at infinity” the technique of arranging decompositions of ternary forms along lines. More precisely, let $r_{\max}(n, d)$ denote the desired answer to the mentioned Waring problem. Elementary considerations show that for each fixed n , $r_{\max}(n, d) = O(d^{n-1})$, and if $r_{\max}(n, d) = c_n d^{n-1} + O(d^{n-2})$ for some constant c_n (as it is reasonable to expect), then it must be $1/n! \leq c_n \leq 1/(n-1)!$. The best general upper bound on $r_{\max}(n, d)$ to our knowledge is given by [1, Corollary 9]. This implies that the constant c_n (if it exists) is at most $2/n!$. Using [5, Proposition 4.1] (see also [2, Theorem 7], [4, Theorem 1]), we deduce $r_{\max}(3, d) \geq \lfloor (d+1)^2/4 \rfloor$. Hence, it must be $1/4 \leq c_3 \leq 1/3$. In the present work, for all ternary forms of degree d we obtain power sum decompositions by considering $\lfloor (d^2 + 6d + 1)/4 \rfloor$ points arranged along d lines. Hence, we have $r_{\max}(3, d) = d^2/4 + O(d)$, that is, $c_3 = 1/4$.

The upper bound we are proving lowers the general upper bound [1, Corollary 9] in the special case $n = 3$ and for $d \geq 6$. Nevertheless, it is not the best we can achieve because our purpose here was to determine the asymptotic leading term as simply as we could. To explain how the method works and why the resulting bound can ulteriorly be lowered, let us consider what happens for a ternary quartic f . For introductory purposes, we now use a geometric language; the technical heart of the paper will be elementary linear algebraic instead. We view our quartic as a point $\langle f \rangle$ in the 14-dimensional projective space of all quartic forms, where fourth powers make a degree 16 Veronese surface. That surface is isomorphic to a plane via quadruple embedding, and exploiting apolarity we get four lines, which embed as rational normal quartics. The four curves are chosen so that their span contains $\langle f \rangle$, but no three of them do the same. Then, by means of successive projections and liftings we get a sequence of essentially binary forms that easily handle power sum decompositions. More precisely, we successively consider decompositions of binary forms of degrees 1, 2, 3, 4, with respective lengths 2, 2, 3, 3. Thus $\text{rk } f \leq 10$. This bound is rather relaxed since $r_{\max}(3, 4) = 7$. Note, however, that generic ranks of binary forms of degrees 1, 2, 3, 4 are 1, 2, 2, 3. Moreover, if one uses [3, Proposition 2.7] instead of Proposition 3.1 here, one gets three (or fewer) lines instead of four. With three lines, the binary forms involved are of degrees 2, 3, 4, and the corresponding generic ranks are 2, 2, 3. This way, with a few additional technical cautions, we can reach the value of $r_{\max}(3, 4)$. Similarly, we can reach $r_{\max}(3, 5) = 10$ in a simpler way than in [8]. For ternary sextics and septic, it is reasonable to expect that the bounds $r_{\max}(3, 6) \leq 14$, $r_{\max}(3, 7) \leq 18$ can be proved with a more or less straightforward extension of the method. However, in the present work we prefer not to set up in detail these results about low-degree forms because there are also reasons to believe that to reach $r_{\max}(3, d)$, further considerations could be in order (maybe an enhanced choice of the lines, if not a completely different strategy). We now outline what these reasons are.

When the present paper was in preparation, a log cabin patchwork like the following was shown to us ⁽¹⁾:




FIGURE 1.

The area of the patches, starting from the center, makes a sequence

$$1, 2, 2, 2, 3, 3, 4, 4, \dots$$

The partial sums are

$$1, 3, 5, 7, 10, 13, 17, 21, \dots$$

¹As strange as it seems, during the lunch break on October 7, 2015, the TV was on and at a certain point the patchwork was shown as a tutorial about sewing in the program “Detto Fatto”, broadcast by the national Italian channel RAI 2.

The first five partial sums agree with the values of $r_{\max}(3, d)$, $d = 1, \dots, 5$ that are known at the time of writing. The picture also clearly shows that the area is asymptotically $d^2/4$. This suggests that $r_{\max}(3, d)$ could be $\lfloor (d^2 + 2d + 5)/4 \rfloor$ for $d \geq 2$. This would mean that [4, Theorem 1] is the best that one can achieve for $n = 3$ and odd $d \geq 3$, and that for even degrees one should be able to raise by one the rank reached by monomials (like [4, Theorem 1] for odd degrees).

The picture suggests how to build sets of points that may give rise to forms with the desired rank. On the other hand, at the moment we do not know how our technique for upper bounds could be improved. To decide for the values $12 \leq r_{\max}(3, 6) \leq 14$ and $17 \leq r_{\max}(3, 7) \leq 18$, would indicate whether or not the “patchwork conjecture” is more promising than a likewise straightforward application of the method of the present article. In any case, we acknowledge that the patchwork helped us recognize that, for the purposes of the present work, to consider d lines (Proposition 3.1) makes things simpler than considering $d - 1$ lines ([3, Proposition 2.7]).

2. PREPARATION

We work over an algebraically closed field \mathbb{K} of characteristic zero and fix two symmetric \mathbb{K} -algebras $S^\bullet = \text{Sym}^\bullet S^1$, $S_\bullet = \text{Sym}^\bullet S_1$; we shall keep this notation throughout the paper. We also assume that an *apolarity pairing* between S^\bullet, S_\bullet is given. It is naturally induced by a perfect pairing $S^1 \times S_1 \rightarrow \mathbb{K}$ (for more details see [7, Introduction]). This amounts to say that S^\bullet, S_\bullet are rings of polynomials in a finite and the same number of indeterminates, acting on each other by constant coefficients partial differentiation. For each $x \in S^\bullet$ and $f \in S_\bullet$ we shall denote by $\partial_x f$ the apolarity action of x on f . For each form (homogeneous polynomial) $f \in S_{d+\delta}$, we shall denote by $f_{\delta,d}$ the *partial polarization* map $S^\delta \rightarrow S_d$ defined by $f_{\delta,d}(x) := \partial_x f$. The *apolar ideal* of $f \in S_d$ is the set of all $x \in S^\bullet$ such that $\partial_x f = 0$. We also define the evaluation of a homogeneous form $x \in S^d$ on a linear form $v \in S_1$, by setting

$$x(v) := \frac{\partial_x v^d}{d!}.$$

The (Waring) *rank* of $f \in S_d$, $d > 0$, denoted by $\text{rk } f$, is the least of the numbers r such that f can be written as a sum of r d th powers of forms in S_1 ⁽²⁾; $r_{\max}(n, d)$ is the maximum of the ranks of all such f when $\dim S_1 = n$. The span of v_1, \dots, v_r in some vector space V will be denoted by $\langle v_1, \dots, v_r \rangle$, and the projective space made of all one-dimensional subspaces $\langle v \rangle \subseteq V$, $v \neq 0$, will be denoted by $\mathbb{P}V$. A *morphism of projective spaces* $\mathbb{P}\varphi : \mathbb{P}V \setminus \mathbb{P}\text{Ker } \varphi \rightarrow \mathbb{P}W$ is a map determined by a linear map $\varphi : V \rightarrow W$ by setting $\mathbb{P}\varphi(\langle v \rangle) := \langle \varphi(v) \rangle$. The sign \perp will refer to orthogonality with respect to the apolarity pairing $S^d \times S_d \rightarrow \mathbb{K}$, when some degree d is fixed (sometimes implicitly).

In [9, Sec. 1.3], building on classical results due to Sylvester, the authors deal with binary forms (i.e. $\dim S_1 = 2$, in our notation). They show that power sum

²Since we are assuming that \mathbb{K} is algebraically closed, when $d > 0$ a form $f \in S_d$ is a sum of r d th powers of linear forms if and only if it is a linear combination of r d th powers of linear forms. Using linear combinations allows one to define Waring rank in degree 0 as well, and of course it would be 1 for every nonzero constant. We prefer not to decide here whether the rank of a nonzero constant should be 1 or be left undefined.

decompositions are closely related with the initial degree of the (homogeneous) apolar ideal, that is, the least degree of a nonzero homogeneous element of that ideal. That is the notion of *length* of a binary form (see [9, Def. 1.32 and Lemma 1.33]), which can be generalized in various ways for forms in more indeterminates: see [9, Def. 5.66]. Nowadays, terms related to length are replaced by similar terms related with rank, probably because of the renewed interest in the interplay with the rank of tensors. In the present paper we need that notion only when the form is essentially binary, and what we really use is only its algebraic property of being the initial degree of the apolar ideal in a ring of binary forms. Note that a form $f \in S_d$ belongs to some subring $T_\bullet = \text{Sym}^\bullet T_1$ with $\dim T_1 = 2$, if and only if $\text{Ker } f_{1,d-1}$ has codimension at most 2 in S^1 (it suffices to take a two-dimensional $T_1 \supseteq \text{Ker } f_{1,d-1}^\perp$). Moreover, f belongs to more than one of such subrings if and only if $\text{Ker } f_{1,d-1}$ has codimension at most 1, in which case the initial degree of the apolar ideal of f in each of the subrings T_\bullet , whatever dual ring T^\bullet one chooses, is always the same (and equal to the codimension). This allows us to state the following definition.

Definition 2.1. Let $f \in S_d$. If f belongs to some ring $T_\bullet = \text{Sym}^\bullet T_1$, contained or containing S_\bullet (with the ring structures preserved), with $\dim T_1 = 2$, then we define the *binary length* of f as the initial degree of its apolar ideal, and we denote it by $\text{bl } f$.

The following definition is also useful.

Definition 2.2. Given $x \in S^\bullet$ and $f \in S_\bullet$, an *x-antiderivative* of f is a polynomial $F \in S_\bullet$ such that $\partial_x F = f$, and when x, f are nonzero and homogeneous we sometimes also say that $\langle F \rangle$ is a $\langle x \rangle$ -*antiderivative* of $\langle f \rangle$. Moreover, still in the homogeneous case $x \in S^\delta$, $f \in S_d$, if a decomposition

$$(1) \quad f = \lambda_1 v_1^d + \cdots + \lambda_r v_r^d, \quad \lambda_1, \dots, \lambda_r \in \mathbb{K}, \quad v_1, \dots, v_r \in S_1,$$

is given and x vanishes on no one of v_1, \dots, v_r , then we define *the x-antiderivative of f* ⁽³⁾ *relative to (1)* as the form

$$F := \frac{d! \lambda_1}{(d+\delta)! x(v_1)} v_1^{d+\delta} + \cdots + \frac{d! \lambda_r}{(d+\delta)! x(v_r)} v_r^{d+\delta};$$

when the powers v_1^d, \dots, v_r^d are linearly independent we also say that the above *x-antiderivative* is *relative to* v_1, \dots, v_r .

Note that, in the above notation, the *x-antiderivative* relative to v_1, \dots, v_r is the unique *x-antiderivative* of f that lies in $\langle v_1^{d+\delta}, \dots, v_r^{d+\delta} \rangle$.

Now we explicitly point out two basic facts that are probably well-known, but for which we are not aware of a direct reference ⁽⁴⁾.

Remark 2.3. Let $h \in S^d$, $h' \in S^{d'}$ and $e \geq d$. Then h' divides h if and only if $S^e \cap \text{Ker } \partial_{h'} \subseteq S^e \cap \text{Ker } \partial_h$. One implication immediately follows from $\partial_{ph'} = \partial_p \circ \partial_{h'}$, and by the same reason we have $S^e \cap \text{Ker } \partial_{h'} \subseteq (h' S^{e-d'})^\perp$. When $h' \neq 0$, since $\partial_{h'}$ maps S^e onto $S^{e-d'}$ we have $\dim S^e \cap \text{Ker } \partial_{h'} = \dim S^e - \dim S^{e-d'}$. Since the

³Or also the $\langle x \rangle$ -antiderivative of $\langle f \rangle$, if $f \neq 0$.

⁴Basic facts like these are heavily scattered in the literature, and we may easily overlook some reference. For instance, [8, Remark 3.6] could have been avoided by using the original reference [6, Theorem 2] instead of [10, 9.2.2.1]. We take this occasion for apologizing for that.

apolarity pairing is nondegenerate in fixed degree, we also have $\dim \left(h' S^{e-d'} \right)^\perp = \dim S^e - \dim S^{e-d'}$ when $h' \neq 0$. Hence $S^e \cap \text{Ker } \partial_{h'} = \left(h' S^{e-d'} \right)^\perp$ (even when $h' = 0$) and, similarly, $S^e \cap \text{Ker } \partial_h = \left(h S^{e-d} \right)^\perp$. Now, to show the converse implication, let us suppose that $S^e \cap \text{Ker } \partial_{h'} \subseteq S^e \cap \text{Ker } \partial_h$, that is, $\left(h' S^{e-d'} \right)^\perp \subseteq \left(h S^{e-d} \right)^\perp$. Again because apolarity is nondegenerate in fixed degree, we deduce that $h S^{e-d} \subseteq h' S^{e-d'}$. Choosing $x \in S^1$ that does not divide h' (we can assume $h' \neq 0$ and $\dim S_1 \geq 2$, since the proof is trivial in the opposite case), we have that h' divides $x^{e-d}h$, hence h' divides h .

Remark 2.4. Let $f \in S^d$ with $d \geq 3$ and $\dim S_1 = 3$. Then f is a d th power if and only if for each $x \in S^1$, $\partial_x f$ is a $(d-1)$ th power. One implication is immediate. Conversely, suppose that for each $x \in S^1$, $\partial_x f$ is a $(d-1)$ th power of some linear form. Let $x, y \in S^1$ be linearly independent. According to [8, Lemma 4.1] we have that $\partial_l f = 0$ for some nonzero $l \in \langle x, y \rangle$ ⁽⁵⁾. Repeating the argument with another two-dimensional subspace of S^1 that does not contain l , we get $m \in S^1$ such that $\partial_m f = 0$ and with l, m linearly independent. Hence $\text{Ker } f_{1,d-1} \geq 2$, and since $\dim S_1 = 3$ we have that f is a d th power of a linear form.

Moreover, let $f \in S_d$, $d \geq 3$, be not a d th power. The set of all $\langle v^{d-1} \rangle$ with $\langle v \rangle \in \mathbb{P}S_1$ is an algebraic (Veronese) variety in $\mathbb{P}S_{d-1}$, and $\mathbb{P}f_{1,d-1}$ is an algebraic morphism from a nonempty Zariski open subset of $\mathbb{P}S^1$ to $\mathbb{P}S_{d-1}$. Then $\partial_x f$ is not a $(d-1)$ th power for all $\langle x \rangle$ in a nonempty open Zariski subset of $\mathbb{P}S^1$.

We end this section with two technical lemmas.

Lemma 2.5. *Let $\langle f \rangle \in \mathbb{P}S_d$ and $\langle x \rangle \in \mathbb{P}S^1$, with $\dim S_1 = 2$. Let I be the apolar ideal of f and set $\ell := \text{bl } f$, $\ell' := d + 2 - \ell$,*

$$(2) \quad W := S_{d+1} \cap \partial_x^{-1}(\langle f \rangle), \quad H := S^{\ell'} \cap I, \quad K := S^{\ell'} \cap xI.$$

Finally, let X be the locus of all $\langle h \rangle \in \mathbb{P}H$ such that h is not squarefree and set $\langle v_\infty \rangle := \langle x \rangle^\perp$, so that

$$\langle v_\infty^{d+1} \rangle = S_{d+1} \cap \text{Ker } \partial_x \subset W.$$

Then

- *there exists an epimorphism of projective spaces*

$$\omega : \mathbb{P}H \setminus \mathbb{P}K \rightarrow \mathbb{P}W, \quad \omega(\langle h \rangle) =: \langle w_h \rangle,$$

such that $\partial_h w_h = 0$ for all $\langle h \rangle$;

- *for all $\langle w \rangle \in \mathbb{P}W \setminus \langle v_\infty^{d+1} \rangle$ but at most one, we have*

$$\text{bl } w = \min \{ \ell + 1, \ell' \};$$

- $X \subsetneq \mathbb{P}H$;
- *for each projective line $\mathbb{P}L \subseteq \mathbb{P}H$ that does not meet $\mathbb{P}K$, the restriction $\mathbb{P}L \rightarrow \mathbb{P}W$ of ω is an isomorphism of projective spaces, and if the line $\mathbb{P}L$ is not contained in X then there exists a cofinite subset $U \subset \mathbb{P}L$ such that for each $\langle h \rangle \in U$ we have*
 - *h has distinct roots $\langle v_1 \rangle, \dots, \langle v_{\ell'} \rangle \in \mathbb{P}S_1$;*

⁵In the statement of the cited result, by mistake, one finds $f \in S_d$ instead $f \in S_{d+1}$ and the condition $d \geq 2$ is omitted; cf. [8, Rem. 2.2].

- $f \in \langle v_1^d, \dots, v_{\ell'}^d \rangle$;
- x vanishes on no one of $v_1, \dots, v_{\ell'}$ and $\langle w_h \rangle$ is the $\langle x \rangle$ -antiderivative of $\langle f \rangle$ relative to $v_1, \dots, v_{\ell'}$.

Proof. For each $h \in H$ and $w \in W$ we have $\partial_{xh}w = 0$, because $\partial_x w \in \langle f \rangle$ and $h \in I$; hence $\partial_h w \in S_{\ell-1} \cap \text{Ker } \partial_x = \langle v_\infty^{\ell-1} \rangle$. Thus we have a bilinear map

$$\beta : H \times W \rightarrow \langle v_\infty^{\ell-1} \rangle, \quad \beta(h, w) := \partial_h w.$$

If $h \in K$, then $h = xh'$ for some $h' \in I$; hence for all $w \in W$ we have $\partial_h w = \partial_{h'} \partial_x w = 0$, because $\partial_x w \in \langle f \rangle$. This shows that K is contained in the left kernel of β . Conversely, if h is in the left kernel, then ∂_h vanishes on W , and in particular on $\langle v_\infty^{d+1} \rangle \subset W$. Hence $h = xh'$ for some $h' \in S^d$, by Remark 2.3 ⁽⁶⁾. Choosing an x -antiderivative w of f , we have $0 = \partial_h w = \partial_{h'} f$, and thus $h' \in I$. We conclude that K is the left kernel of β .

Let

$$\bar{\beta} : H \rightarrow \text{Hom}(W, \langle v_\infty^{\ell-1} \rangle), \quad \bar{\beta}(h)(w) := \beta(h, w) = \partial_h w,$$

be the homomorphism induced by β , and let $\iota : W \rightarrow \text{Hom}(W, \langle v_\infty^{\ell-1} \rangle)$ be an isomorphism such that $\iota(w)(w) = 0$ for all $w \in W$ (in other words, ι is the homomorphism induced by a nondegenerate bilinear alternating map on W with values in $\langle v_\infty^{\ell-1} \rangle$, which certainly exists because $\dim W = 2$). Then $\varphi := \iota^{-1} \circ \bar{\beta} : H \rightarrow W$ is a linear map with kernel K such that $\partial_h(\varphi(h)) = 0$ for all $h \in H$. This shows that $\omega := \mathbb{P}\varphi$ is a morphism of projective spaces such that $\partial_h w_h = 0$ (under the notation $\langle w_h \rangle := \omega(\langle h \rangle) = \langle \varphi(h) \rangle$). We have to check that ω is surjective.

According to [9, Theorem 1.44(iv)], I is generated by two homogeneous forms $l \in S^\ell, h^0 \in S^{\ell'}$ (hence $h^0 \in H$). Recall also that $\ell \leq \ell'$ because $\ell = \text{bl } f$. Therefore

$$(3) \quad H = lS^{\ell'-\ell} + \langle h^0 \rangle, \quad K = lS^{\ell'-\ell-1}.$$

Since $S^{d+1} \subset I$, we have that l, h^0 are coprime, and therefore $h^0 \notin lS^{\ell'-\ell}$. Since $\dim \varphi(lS^{\ell'-\ell}) = 1$ we have that φ is surjective, and hence ω is surjective as it was to be shown.

Let $\varphi(lS^{\ell'-\ell}) =: \langle w_l \rangle \in \mathbb{P}W$ (possibly $\langle w_l \rangle = \langle v_\infty^{d+1} \rangle$). Since $\partial_l w_l = 0$ for all $p \in S^{\ell'-\ell}$, we have $\partial_l w_l = 0$. Note also that $w_l = w_{pl}$ for all $p \in S^{\ell'-\ell} \setminus xS^{\ell'-\ell-1}$. Moreover,

$$(4) \quad \langle h \rangle, \langle h' \rangle \in \mathbb{P}H \setminus \mathbb{P}K, \langle w_{h'} \rangle \neq \langle w_h \rangle \implies \partial_h w_{h'} \neq 0,$$

because $\partial_h w_{h'} = 0$ would imply that ∂_h vanishes on $\langle w_h, w_{h'} \rangle = W$ ($\dim W = 2$), and this is excluded since $h \notin K$. Since ω is surjective, we conclude that $\partial_l w \neq 0$ for each $\langle w \rangle \in \mathbb{P}W \setminus \{\langle w_l \rangle\}$. On the other hand, if $\langle w \rangle \in \mathbb{P}W \setminus \{\langle v_\infty^{d+1} \rangle\}$, then $\langle \partial_x w \rangle = \langle f \rangle$, and hence the apolar ideal of w is contained in I and contains xI . Thus

$$\ell \leq \text{bl } w \leq \ell + 1, \quad \forall \langle w \rangle \in \mathbb{P}W \setminus \{\langle v_\infty^{d+1} \rangle\}.$$

Now, if $\ell' \geq \ell + 1$, then for each $\langle w \rangle \in \mathbb{P}W \setminus \{\langle v_\infty^{d+1} \rangle, \langle w_l \rangle\}$ we have $\text{bl } w = \ell + 1 = \min\{\ell + 1, \ell'\}$. To deal with the case $\ell' = \ell$, notice that for each $w \in W$ we

⁶Alternatively, one may observe that $0 = \partial_h(v_\infty^{d+1}) = \frac{(d+1)!}{\ell'!} h(v_\infty) v_\infty^{\ell-1}$. Hence h vanishes on the root v_∞ of x , that is, h is divisible by x .

have $\partial_h w = 0$ for some $\langle h \rangle \in H$, because φ is surjective; hence $\text{bl } w \leq \ell'$. Thus, if $\ell = \ell'$ then for each $\langle w \rangle \in \mathbb{P}W \setminus \langle v_\infty^{d+1} \rangle$ we have $\text{bl } w = \ell' = \min \{\ell + 1, \ell'\}$ ⁽⁷⁾.

Since l, h^0 are coprime, taking into account (3) and Bertini's theorem (see also [11, Lemma 1.1, Remark 1.1.1]), we have that X is a proper subset of $\mathbb{P}H$.

Finally, let $\mathbb{P}L \subseteq \mathbb{P}H \setminus \mathbb{P}K$ be a projective line. The restriction $\mathbb{P}L \rightarrow \mathbb{P}W$ of ω is an isomorphism simply because $\mathbb{P}W$ is a projective line as well, and $\mathbb{P}L \cap \mathbb{P}K = \emptyset$. Since the proper subset $X \subsetneq \mathbb{P}H$ is algebraic, with equation given by the discriminant of degree ℓ' forms (inside $\mathbb{P}H$), we have that if $\mathbb{P}L \not\subseteq X$, then $U := \mathbb{P}L \setminus (X \cup \omega^{-1}(\langle v_\infty^{d+1} \rangle))$ is a cofinite subset of $\mathbb{P}L$. Since each $\langle h \rangle \in U$ is outside X , h is squarefree, that is, it has distinct roots $\langle v_1 \rangle, \dots, \langle v_{\ell'} \rangle \in \mathbb{P}S_1$. For such $h, v_1, \dots, v_{\ell'}$, according to [9, Lemma 1.31], we have $f \in \langle v_1^d, \dots, v_{\ell'}^d \rangle$ as required. By the same reason, we have $w_h \in \langle v_1^{d+1}, \dots, v_{\ell'}^{d+1} \rangle$, and since $\langle w_h \rangle \neq \langle v_\infty^{d+1} \rangle$, $\langle w_h \rangle$ is a $\langle x \rangle$ -antiderivative of $\langle f \rangle$. Moreover, x vanishes on no one of $v_1, \dots, v_{\ell'}$ by (4), and $v_1^d, \dots, v_{\ell'}^d$ are linearly independent because $\ell' \leq d + 1$. The above said suffices to prove that $\langle w_h \rangle$ is the $\langle x \rangle$ -antiderivative of $\langle f \rangle$ relative to $v_1, \dots, v_{\ell'}$. \square

Lemma 2.6. *Let $\langle g' \rangle \in \mathbb{P}S_d$ with $\dim S_1 = 3$, $0 < d = 2s + \varepsilon$, $\varepsilon \in \{0, 1\}$ and s integer. Let $\langle l^0 \rangle, \dots, \langle l^t \rangle \in \mathbb{P}S^1$ be distinct and such that $\partial_{l^0} g' = 0$, and for each $i \in \{1, \dots, t\}$ let g_i be an l^i -antiderivative of g' . If*

$$\text{bl } g' = \text{bl } \partial_{l^0} g_1 = \dots = \text{bl } \partial_{l^0} g_t = s + 1$$

(⁸), then there exists a power sum decomposition

$$(5) \quad g' = v_1^d + \dots + v_r^d$$

such that: $r \leq s + 1 + \varepsilon$ and, for each $i \in \{1, \dots, t\}$,

- l^i vanishes on no one of v_1, \dots, v_r ,
- denoting by F_i the l^i -antiderivative relative to (5), $\text{bl } (g_i - F_i) = s + 1 + \varepsilon$.

Proof. For each $i \in \{0, \dots, t\}$, let $R_i^\bullet := S^\bullet / (l^i)$, $R_{i,\bullet} := \text{Ker } \partial_{l^i} \subset S_\bullet$, with the apolarity pairing induced by the one between S^\bullet and S_\bullet . Let $I \subset R_0^\bullet$ be the apolar ideal of $g' \in R_{0,d}$, set

$$H := R_0^{s+1+\varepsilon} \cap I,$$

for each $i \in \{1, \dots, t\}$ set

$$W_{0,i} := R_{0,d+1} \cap \partial_{l^i}^{-1}(\langle g' \rangle)$$

and when $\varepsilon = 1$, also $\langle k_i \rangle := R_0^{s+2} \cap l^i I$. For each $i \in \{1, \dots, t\}$, let us exploit Lemma 2.5 with R_0^\bullet , $R_{0,\bullet}$, g' , $l^i + (l^0)$ in place of S^\bullet , S_\bullet , f , x . We get projective epimorphisms

$$\omega_i : \mathbb{P}H \rightarrow \mathbb{P}W_{0,i}.$$

Moreover, we can fix a projective line $\mathbb{P}L \subseteq \mathbb{P}H$ ($\mathbb{P}L = \mathbb{P}H$ when $\varepsilon = 0$) not contained in the singular locus X (which does not depend on i) and passing through no one of $\langle k_1 \rangle, \dots, \langle k_t \rangle$ (when $\varepsilon = 1$). Hence, the restriction $\varrho_i : \mathbb{P}L \rightarrow \mathbb{P}W_{0,i}$ of ω_i is a projective isomorphism for each i , and we also have cofinite subsets $U_{0,i} \subset \mathbb{P}L$ that fulfill the properties listed by the end of the statement of Lemma 2.5.

⁷The equality $\text{bl } w = \min \{\ell + 1, \ell'\}$ we have just proved for all $\langle w \rangle \in \mathbb{P}W \setminus \{\langle v_\infty^{d+1} \rangle, \langle w_l \rangle\}$ says, in other terms, that $\text{bl } w = \ell + 1$ unless d is even, $d = 2s$, and ℓ is the maximum allowed for that degree, that is, $s + 1$.

⁸For each i , $\partial_{l^0} g_i$ is annihilated by l^i .

Now, for each i we have $\partial_{l^0} g_i \neq 0$ because $\text{bl } \partial_{l^0} g_i = s+1 > 0$. Hence the vector space

$$W_{i,0} := R_{i,d+1} \cap \partial_{l^0}^{-1}(\langle \partial_{l^0} g_i \rangle)$$

is two-dimensional. Since $W_{0,i} = R_{0,d+1} \cap \partial_{l^i}^{-1}(\langle g' \rangle)$, for all $w \in W_{0,i}$ we have

$$\partial_{l^i} w = \lambda_i(w) g'$$

for some scalar $\lambda_i(w)$, and therefore $\lambda_i(w) g_i - w \in W_{i,0}$. This defines a map $W_{0,i} \rightarrow W_{i,0}$ and to check that it is a vector space isomorphism is easy (take into account that $\partial_{l^0}(\lambda_i(w) g_i - w) = \lambda_i(w) \partial_{l^0} g_i$). Therefore we have isomorphisms of projective spaces

$$\tau_i : \mathbb{P}W_{0,i} \rightarrow \mathbb{P}W_{i,0}, \quad \langle w \rangle \mapsto \langle \lambda_i(w) g_i - w \rangle.$$

According to Lemma 2.5, we have cofinite subsets $U'_{i,0} \subset \mathbb{P}W_{i,0}$ such that

$$(6) \quad \text{bl } w = s+1+\epsilon, \quad \forall \langle w \rangle \in U'_{i,0}$$

(more precisely, $\#(\mathbb{P}W_{i,0} \setminus U'_{i,0}) \leq 2$).

Let $U_{i,0} := \varrho_i^{-1}(\tau_i^{-1}(U'_{i,0}))$ for each i , which is obviously a cofinite subset of $\mathbb{P}L$. Now, let us pick $\langle h \rangle$ in the nonempty intersection

$$U_{0,1} \cap \dots \cap U_{0,t} \cap U_{1,0} \cap \dots \cap U_{t,0},$$

and let $\langle v_1 \rangle, \dots, \langle v_{s+1+\epsilon} \rangle$ be its roots, which are distinct because $\langle h \rangle \in U_{0,i}$ (whatever i one chooses). For each i , l^i vanishes on no one of $v_1, \dots, v_{s+1+\epsilon}$, because $\langle h \rangle \in U_{0,i}$. Since $g' \in \langle v_1^d, \dots, v_{s+1+\epsilon}^d \rangle$, $d > 0$, for an appropriate choice of the representatives $v_1, \dots, v_{s+1+\epsilon}$ one gets (5). Since F_i is the l^i -antiderivative of g' relative to (5), that is, relative to $v_1, \dots, v_{s+1+\epsilon}$, we have $\langle F_i \rangle = \omega_i(\langle h \rangle)$. Since F_i is an l^i -antiderivative of g' , we have $\lambda_i(F_i) = 1$, and hence

$$\tau_i(\omega_i(\langle h \rangle)) = \langle g_i - F_i \rangle.$$

Since $\langle h \rangle \in U_{i,0}$ for each i , we have $\langle g_i - F_i \rangle \in U'_{i,0}$, and therefore $\text{bl}(g_i - F_i) = s + \epsilon + 1$ by (6). \square

3. THE UPPER BOUND

Proposition 3.1. *Let $f \in S_d$, with $\dim S_1 = 3$ and $d \geq 2$, let $f_1, \dots, f_a \in S_\bullet$ be homogeneous forms with degrees at least $d+1$, and let $\langle x^1 \rangle, \dots, \langle x^b \rangle \in \mathbb{P}S^1$. If f, f_1, \dots, f_a are not powers of linear forms, then there exist distinct*

$$\langle l^1 \rangle, \dots, \langle l^d \rangle \in \mathbb{P}S^1 \setminus \{\langle x^1 \rangle, \dots, \langle x^b \rangle\}$$

such that

$$\partial_{l^1 \dots l^d} f = 0, \quad \partial_{l^1 \dots \hat{l}^i \dots l^d} f \neq 0 \quad \forall i \in \{1, \dots, d\}, \quad \partial_{l^1 \dots l^d} f_j \neq 0 \quad \forall j \in \{1, \dots, a\},$$

where the hat denotes omission.

Proof. According to Remark 2.4, we can fix

$$\langle l^d \rangle \in \mathbb{P}S^1 \setminus \{\langle x^1 \rangle, \dots, \langle x^b \rangle\}$$

such that $f'_1 := \partial_{l^d} f_1, \dots, f'_a := \partial_{l^d} f_a$ are not powers of linear forms. Since $f \neq 0$ because it is not a d th power, we can also assume that, in addition, $f' := \partial_{l^d} f \neq 0$.

Let us first suppose that $d = 2$. For each $i \in \{1, \dots, a\}$, since f'_i is not a power of a linear form, we may have $\partial_{l^i} f'_i = 0$ for at most one $\langle l \rangle \in \mathbb{P}S^1$. Therefore there exists a finite subset $\Sigma \subset \mathbb{P}S^1$ such that for all $\langle l \rangle \in \mathbb{P}S^1 \setminus \Sigma$ and $i \in \{1, \dots, a\}$

we have $\partial_l f'_i \neq 0$. We have $\partial_l f = 0$ for at most one $\langle l \rangle \in \mathbb{P}S^1$ as well. Since $\text{Ker } f'_{1,0} = \langle f' \rangle^\perp$ is an infinite set, we can pick out $\langle l^1 \rangle \in \mathbb{P} \langle f' \rangle^\perp$ such that

- $\partial_{l^1} f \neq 0$;
- $\langle l^1 \rangle \notin \{ \langle l^2 \rangle, \langle x^1 \rangle, \dots, \langle x^b \rangle \} \cup \Sigma$.

It is immediate to check that $\langle l^1 \rangle, \langle l^2 \rangle$ fulfill all the requirements in the statement.

Now, let us assume $d \geq 3$. In this case we can assume that f' is not a $(d-1)$ th power and, by induction on d , that the statement holds with f' in place of f , with f, f'_1, \dots, f'_a in place of f_1, \dots, f_a and with l^d, x^1, \dots, x^b in place of x^1, \dots, x^b . This gives linear forms l^1, \dots, l^{d-1} that, together with l^d , fulfill all the requirements. \square

Proposition 3.2. *Let $f \in S_d$ with $\dim S_1 = 3$, $e \in \{0, \dots, d\}$, $e = 2s + \varepsilon$, with $\varepsilon \in \{0, 1\}$ and s integer, and let*

$$\langle l^1 \rangle, \dots, \langle l^d \rangle \in \mathbb{P}S^1$$

be distinct and such that

$$\partial_{l^1 \dots l^d} f = 0; \quad \partial_{l^1 \dots \widehat{l^i} \dots l^d} f \neq 0 \ \forall i.$$

Then there exists a power sum decomposition

$$(7) \quad \partial_{l^{e+1} \dots l^d} f = v_1^e + \dots + v_r^e$$

such that:

- $r \leq s^2 + 3s + \varepsilon(s+2)$;
- for each $i \in \{e+1, \dots, d\}$, l^i vanishes on no one of v_1, \dots, v_r and denoting by F_i the l^i -antiderivative relative to (7), we have

$$\text{bl}(\partial_{l^{e+1} \dots \widehat{l^i} \dots l^d} f - F_i) = s+1 + \varepsilon.$$

Proof. When $e = 0$ it suffices to define (7) as the decomposition of 0 with no summands. By induction, we can assume that $e \geq 1$ and that the proposition holds with $e-1$ in place of e . Therefore we get a decomposition

$$(8) \quad \partial_{l^e \dots l^d} f = v_1'^{e-1} + \dots + v_{r'}'^{e-1}$$

such that

$$(9) \quad r' \leq s^2 + 2s - 1 + \varepsilon(s+1)$$

and each of l^e, \dots, l^d vanishes on no one of $v_1', \dots, v_{r'}'$. We can also consider for each $i \in \{e, \dots, d\}$ the l^i -antiderivative relative to (8), which we denote by G'_i , and set

$$g'_i := \partial_{l^e \dots \widehat{l^i} \dots l^d} f - G'_i,$$

so that

$$\text{bl } g'_i = s+1.$$

For each $i \in \{e+1, \dots, d\}$, let G_i be the $l^e l^i$ -antiderivative relative to (8) and set

$$g_i := \partial_{l^{e+1} \dots \widehat{l^i} \dots l^d} f - G_i,$$

so that

$$\partial_{l^e} g_i = g'_i, \quad \partial_{l^i} g_i = g'_e.$$

By the above construction, we can exploit Lemma 2.6 with $e, g'_e, l^e, \dots, l^d, g_{e+1}, \dots, g_d$ in place of $d, g', l^0, \dots, l^t, g_1, \dots, g_t$. We get a decomposition

$$(10) \quad g'_e = v_1^e + \dots + v_{r''}^e$$

such that

$$(11) \quad r'' \leq s + 1 + \varepsilon ,$$

each of l^{e+1}, \dots, l^d vanishes on no one of $v_1, \dots, v_{r''}$ and denoting by H_i the l^i -antiderivative relative to (10), we have

$$(12) \quad \text{bl}(g_i - H_i) = s + 1 + \varepsilon .$$

Since we defined G'_e as the l^e -antiderivative relative to (8), by taking suitable multiples of $v'_1, \dots, v'_{r'}$ and calling them $v_{r''+1}, \dots, v_{r''+r'}$, respectively, we have

$$G'_e = v_{r''+1}^e + \dots + v_r^e ,$$

with $r := r'' + r'$. By definition of g'_e and by (10) we conclude that

$$\partial_{l^{e+1} \dots l^d} f = g'_e + G'_e = v_1^e + \dots + v_r^e .$$

To show that the above is the required decomposition (7), first note that (9) and (11) give

$$r \leq s^2 + 3s + \varepsilon(s + 2) ,$$

as it was to be shown. Moreover, since each of l^e, \dots, l^d vanishes on no one of $v'_1, \dots, v'_{r'}$, which are proportional to $v_{r''+1}, \dots, v_r$, and each of l^{e+1}, \dots, l^d vanishes on no one of $v_1, \dots, v_{r''}$, we have that for each $i \in \{e + 1, \dots, d\}$, l^i vanishes on no one of v_1, \dots, v_r . Finally, for the l^i -antiderivatives F_i s we have

$$F_i = H_i + G_i = H_i + \partial_{l^{e+1} \dots \widehat{l^i} \dots l^d} f - g_i ,$$

hence

$$\partial_{l^{e+1} \dots \widehat{l^i} \dots l^d} f - F_i = g_i - H_i$$

and the last requirement to be fulfilled follows from (12). \square

Proposition 3.3. *When $\dim S_1 = 3$, $d > 0$, for all $f \in S_d$ we have*

$$\text{rk } f \leq \left\lfloor \frac{d^2 + 6d + 1}{4} \right\rfloor .$$

Proof. If f is a d th power then $\text{rk } f \leq 1$ and the result trivially follows. Hence we can assume that f is not a d th power. Exploiting Proposition 3.1 with $a = b = 0$, we get $\langle l^1 \rangle, \dots, \langle l^d \rangle \in \mathbb{P}S^1$ such that

$$\partial_{l^1 \dots l^d} f = 0 , \quad \partial_{l^1 \dots \widehat{l^i} \dots l^d} f \neq 0 \ \forall i \in \{1, \dots, d\} .$$

Now the result immediately follows from Proposition 3.2 with $e := d$. \square

Proposition 3.4. *We have $r_{\max}(3, d) = d^2/4 + O(d)$.*

Proof. An immediate consequence of Proposition 3.3 together with [5, Proposition 4.1] (see also [2, Theorem 7], [4, Theorem 1]). \square

REFERENCES

- [1] Blekherman, Grigoriy and Teitler, Zach. On maximum, typical and generic ranks. *Math. Ann.*, **362**, No. 3-4 (2015) 1021-1031, DOI: 10.1007/s00208-014-1150-3
- [2] Buczyńska, Weronika and Buczyński, Jarosław and Teitler, Zach. Waring decompositions of monomials. *J. Algebra*, **378** (2013) 45–57, DOI: 10.1016/j.jalgebra.2012.12.011
- [3] E. Ballico and A. De Paris. Generic Power Sum Decompositions and Bounds for the Waring rank. arXiv:1312.3494v1 [math.AG].
- [4] Buczyński, Jarosław and Teitler, Zach. Some examples of forms of high rank. preprint (2015), arXiv:1503.08253 [math.AG].
- [5] Carlini, Enrico and Catalisano Maria V. and Geramita, Anthony V. The solution to the Waring problem for monomials and the sum of coprime monomials. *J. Algebra*, **370** (2012) 5–14.
- [6] Comas, G. and Seiguer, M. On the rank of a binary form. *Found. Comput. Math.*, **11**(1) (2011) 65-78.
- [7] De Paris, Alessandro. A proof that the maximum rank for ternary quartics is seven. *Matematiche (Catania)*, **25**, No. 2 (2015), to appear. cf. arXiv:1309.6475v1 [math.AG].
- [8] De Paris, Alessandro. Every ternary quintic is a sum of ten fifth powers. *Int. J. Algebra Comput.*, **25**, No. 4 (2015) 607–631, DOI: 10.1142/S0218196715500125
- [9] Iarrobino, Anthony and Kanev, Vassil. *Power sums, Gorenstein algebras, and determinantal loci*. Lecture Notes in Mathematics, vol. 1721, Springer-Verlag, Berlin, 1999. Appendix C by Iarrobino and Steven L. Kleiman.
- [10] Landsberg, Joseph M. *Tensors: geometry and applications*. American Mathematical Society (AMS), Providence, RI, 2012.
- [11] Kleppe, Johannes. Representing a homogenous polynomial as a sum of powers of linear forms. Thesis for the degree of Candidatum Scientiarum, Department of Mathematics, Univ. Oslo (1999).
- [12] Teitler, Zach. Geometric Lower Bounds for Generalized Ranks. preprint (2014), arXiv:1406.5145v2 [math.AG].

DIPARTIMENTO DI MATEMATICA E APPLICAZIONI “RENATO CACCIOPOLI”,
 UNIVERSITÀ DI NAPOLI FEDERICO II (ITALY)
E-mail address: `deparis@unina.it`